arXiv:1305.1117v1 [math.AT] 6 May 2013

The T-equivariant Integral Cohomology Ring of F_4/T

Takashi SATO *

Abstract

We determine the T-equivariant integral cohomology of F_4/T combinatorially by the GKM theory, where T is a maximal torus of the exceptional Lie group F_4 and acts on F_4/T by the left multiplication.

1 Introduction and statement of the result

The cohomology of homogeneous spaces are studied by many mathematicians, for example A. Borel and R. Bott. Let G be a simple, compact, connected Lie group and T its maximal torus. The homogeneous space G/T is a flag variety and it plays an important role in topological geometry, algebraic geometry, representation theory, and combinatorics. It is natural to consider the problem to determine the T-equivariant integral cohomology ring $H_T^*(G/T) = H^*(ET \times_T G/T)$, where T acts on G/T by the left multiplication.

Goresky, Kottwitz, and MacPherson [GKM] gave a powerful method to determine the equivariant cohomology with $\mathbb{Q}$ coefficients of some good spaces. It is called the GKM theory. Let us explain how the GKM theory works in our situation. Since the fixed points set $(G/T)^T$ is the Weyl group $W(G)$, the inclusion $i\colon (G/T)^T \to G/T$ induces the map

$$i^*\colon H_T^*(G/T) \to H_T^*((G/T)^T) = \prod_{W(G)} H^*(BT) = \mathrm{Map}(W(G), H^*(BT)).$$

Tensoring with $\mathbb{Q}$, i^* is injective by the localization theorem. Then Guillemin and Zara [GZ] restated an important theorem of the GKM theory as: the T-equivariant cohomology $H_T^*(G/T)$ is completely determined by a graph with additional data obtained from G. Especially they defined the "cohomology" ring of the graph as a subring of $\mathrm{Map}(W(G), H^*(BT))$ and showed it coincides with the image of i^*. This graph is called a GKM graph. Harada, Henriques, and Holm [HHH] showed that i^* is injective with integral coefficients whenever G is not of type C.

As a consequence of concrete computations by the GKM theory, for a simple Lie group G of classical types and type G_2, Fukukawa, Ishida, and Masuda [FIM], [F] determined the cohomology ring of the GKM graph of G/T. They determined the equivariant integral cohomology rings $H_T^*(G/T)$ for a Lie group G of type A, B, D, and G_2. In this paper we determine the T-equivariant integral cohomology ring of F_4/T by the GKM theory. The ordinary integral cohomology ring $H^*(F_4/T)$ was determined by Toda-Watanabe [TW].

---

*`t-sato@math.kyoto-u.ac.jp`

For $x = (x_1, \dots, x_n)$, let $e_i(x)$ denote the i^{th} elementary symmetric polynomial in $x_1, \dots, x_n$. Put $x^k = (x_1^k, \dots, x_n^k)$. For a linear transformation α of $\mathbb{R}x_1 \oplus \cdots \oplus \mathbb{R}x_n$, let $\alpha x = (\alpha x_1, \dots, \alpha x_n)$. Then $e_i(x^k)$ and $e_i(\alpha x)$ denote the i^{th} elementary symmetric polynomial in $x_1^k, \dots, x_n^k$ and $\alpha x_1, \dots, \alpha x_n$, respectively. The following theorem is the main result of this paper. In this theorem $t = (t_1, t_2, t_3, t_4)$, $\tau = (\tau_1, \tau_2, \tau_3, \tau_4)$, and ρ is the linear transformation of $\mathbb{R}t_1 \oplus \cdots \oplus \mathbb{R}t_4$ defined as (3.2).

Theorem 1.1. *Let T be a maximal torus of F_4 which acts on F_4/T by the left multiplication, then the T-equivariant integral cohomology ring of F_4/T is given as:*

$$H_T^*(F_4/T) \cong \mathbb{Z}[t_i, \gamma, \tau_i, \gamma_1, \gamma_3, \omega \mid 1 \le i \le 4]/(r_1', r_1, r_2, r_3, r_4, r_6, r_8, r_{12}),$$

where $|t_i| = |\gamma| = |\tau_i| = |\gamma_1| = 2$, $|\gamma_3| = 6$, and $|\omega| = 8$. Put $A = \gamma_1^4 + \gamma_1^2 e_1(t)e_1(\tau) - \gamma_3 e_1(\tau) - \gamma_1 e_3(t) + \gamma_1^2 e_2(t) + \gamma_1 e_1(t)e_2(t)$, then

$$\begin{aligned}
r_1' &= 2\gamma - e_1(t),\\
r_1 &= 2\gamma_1 - (e_1(\tau) - e_1(t)),\\
r_2 &= 2\gamma_1^2 - (e_2(\tau) - e_2(t)) + (e_1(\tau) - e_1(t))e_1(t),\\
r_3 &= 2\gamma_3 - (e_3(\tau) - e_3(t)),\\
r_4 &= 2A + e_4(\tau) - e_4(t) - 3\omega,\\
r_6 &= \gamma_3^2 + \gamma_3 e_3(t) - \gamma_1^2 e_4(\tau) - \gamma_1 e_1(t)e_4(\tau) - (\omega - A)e_2(t) - \gamma^2\omega,\\
r_8 &= e_4(\tau)^2 - A^2 + (e_4(\tau) + 2\omega)A - e_4(\tau)(2\omega + e_4(t)) - \omega e_4(\rho t),\\
r_{12} &= \omega(\omega - e_4(\rho t))(\omega + e_4(\rho^2 t)).
\end{aligned}$$

We can obtain the integral cohomology ring of F_4/T as a corollary of Theorem 1.1 as follows. There is a fibration sequence

$$F_4/T \longrightarrow ET \times_T F_4/T \xrightarrow{p} BT.$$

Since the projection $p\colon ET \times_T F_4/T \to BT$ restricts to $\pi\colon ET \times_T (F_4/T)^T \to BT$ by composing the inclusion $i\colon ET \times_T (F_4/T)^T \to ET \times_T F_4/T$, the induced map $\pi^*\colon H^*(BT) \to H^*(ET \times_T (F_4/T)^T) = \mathrm{Map}(W(F_4), H^*(BT))$ sends elements of $H^*(BT)$ to constant functions. In Theorem 1.1, t_1, t_2, t_3, t_4, and γ are constant functions (see Section 4). Since the cohomology of F_4/T and BT have vanishing odd parts, the Serre spectral sequence of the fibration p collapses at the E_2-term. Hence $H^*(F_4/T) \cong H_T^*(F_4/T)/(t_1, t_2, t_3, t_4, \gamma)$.

Corollary 1.1 ([TW, Theorem A]). *The integral cohomology ring of F_4/T is given as:*

$$H^*(F_4/T) \cong \mathbb{Z}[\tau_i, \gamma_1, \gamma_3, \omega \mid 1 \le i \le 4]/(\overline{r}_1, \overline{r}_2, \overline{r}_3, \overline{r}_4, \overline{r}_6, \overline{r}_8, \overline{r}_{12}),$$

where

$$\begin{aligned}
\overline{r}_1 &= 2\gamma_1 - e_1(\tau),\\
\overline{r}_2 &= 2\gamma_1^2 - e_2(\tau),\\
\overline{r}_3 &= 2\gamma_3 - e_3(\tau),\\
\overline{r}_4 &= e_4(\tau) - 2\gamma_1 e_3(\tau) + 2\gamma_1^4 - 3\omega,\\
\overline{r}_6 &= -\gamma_1^2 e_4(\tau) + \gamma_3^2,\\
\overline{r}_8 &= 3e_4(\tau)\gamma_1^4 - \gamma_1^8 + 3\omega(\omega + e_3(\tau)\gamma_1),\\
\overline{r}_{12} &= \omega^3.
\end{aligned}$$

Corollary 1.1 will be proved in Section 8. Throughout this paper, all cohomology groups and rings will be taken with integral coefficients.

2 GKM graph and its cohomology

Let G be a compact connected Lie group and let T be its maximal torus. Specializing and abstracting the work of Goresky, Kottwitz, and MacPherson [GKM], Guillemin and Zara [GZ] introduced a certain graph to each of whose edge an element of $H^2(BT)$ is given and showed the T-equivariant cohomology of G/T with complex coefficient is recovered from this graph. Let us introduce this special graph. Recall that there is a natural identification

$$\mathrm{Hom}(T, S^1) \cong H^2(BT),$$

where the left hand side is the set of weights of G. Let $W(G)$ and $\Phi(G)$ denote the Weyl group and the root system of G, respectively. Since every root is a weight, we regard $\Phi(G) \subset H^2(BT)$. There is a canonical action of the Weyl group $W(G)$ on $\mathrm{Hom}(T, S^1)$ and it restricts to $\Phi(G)$. We denote this action as $w\alpha$ for $w \in W(G)$ and $\alpha \in H^2(BT)$. Recall that each $\alpha \in \Phi(G)$, one can assign a reflection σ_α which is an element of the Weyl group $W(G)$.

Definition 2.1. The GKM graph of G/T is the Cayley graph of $W(G)$ with respect to a generating set $\{\sigma_\alpha \in W(G) \mid \alpha \in \Phi(G)\}$ which is equipped with the cohomology classes $\pm w\alpha \in H^2(BT)$ to the edge ww' satisfying $w' = w\sigma_\alpha$. We call $\pm w\alpha$ the label of the edge ww'.

The ambiguity of the sign of the label $\pm w\alpha$ occurs from the equation $w'\alpha = w\sigma_\alpha\alpha = -w\alpha$. Let us introduce the cohomology of a GKM graph. Consider a function $f : W(G) \to H^*(BT)$ between sets. We say that f satisfies the GKM condition or f is a GKM function if for any $w \in W(G)$ and $\alpha \in \Phi(G)$,

$$f(w) - f(w\sigma_\alpha) \in (w\alpha) \subset H^*(BT),$$

where $(x_1, \ldots, x_n)$ means the ideal generated by $x_1, \ldots, x_n$. It is easy to see that all GKM functions form a subring of $\prod_{W(G)} H^*(BT)$, where we identify the set of all functions $W(G) \to H^*(BT)$ with $\prod_{W(G)} H^*(BT)$. Since the GKM graph of G/T has $W(G)$ as its vertex set, a GKM function assigns an element of $H^*(BT)$ to each vertex of the GKM graph.

Definition 2.2. Let $\mathcal{G}$ be the GKM graph of G/T. The cohomology ring $H^*(\mathcal{G})$ is defined as the subring of $\prod_{W(G)} H^*(BT)$ consisting of all GKM functions.

Guillemin and Zara [GZ, Theorem 1.7.3] restated an important theorem of the GKM theory as

$$H_T^*(G/T;\mathbb{C}) \cong H^*(\mathcal{G}) \otimes \mathbb{C}.$$

Harada, Henriques, and Holm refined this result to the integral cohomology. More precisely, we have:

Theorem 2.1 ([HHH, Theorem 3.1 and Lemma 5.2])**.** *Suppose the Lie group G is simple and let $\mathcal{G}$ be the GKM graph of G/T. If G is not of type C_l, then there is an isomorphism*

$$H_T^*(G/T) \cong H^*(\mathcal{G}).$$

3 The GKM graph of F_4/T

In this section we describe and analyze the GKM graph of F_4/T. First of all let us choose a maximal torus of F_4. Let T^4 be the standard maximal torus of $SO(9)$ and let $\overline{t}_1$, $\overline{t}_2$, $\overline{t}_3$, $\overline{t}_4 \in H^2(BT^4)$ be the canonical basis. For the universal covering $\mu : Spin(9) \to SO(9)$ let $T = \mu^{-1}(T^4)$. Then T is a maximal torus of $Spin(9)$. Since $Spin(9)$ is a Lie subgroup of F_4 (cf. [A]), T is also a maximal torus of F_4. We fix a maximal torus of F_4 to T. Let t_i denote $\mu^*(\overline{t}_i) \in H^2(BT)$. By definition we have

$$H^*(BT) = \mathbb{Z}[t_1, t_2, t_3, t_4, \gamma]/(2\gamma - e_1(t)).$$

To describe the Weyl group $W(F_4)$ we start with the root system of F_4. The root system $\Phi(F_4)$ is given as:

$$\Phi(F_4) = \{\pm(t_i + t_j), \pm(t_i - t_j), \pm t_k, \frac{1}{2}(\pm t_1 \pm t_2 \pm t_3 \pm t_4) \mid 1 \le i < j \le 4,\ 1 \le k \le 4\}$$

The roots $\pm(t_i + t_j)$ and $\pm(t_i - t_j)$ are called long roots, and $\pm t_k$ and $\frac{1}{2}(\pm t_1 \pm t_2 \pm t_3 \pm t_4)$ are called short roots. Put

$$\alpha_1 = t_2 - t_3, \qquad \alpha_2 = t_3 - t_4,$$
$$\alpha_3 = t_4, \qquad \alpha_4 = \frac{1}{2}(t_1 - t_2 - t_3 - t_4).$$

Then the Dynkin diagram of F_4 is as:



Then $W(F_4)$ is generated by the reflections σ_{α_i} for $i = 1, 2, 3, 4$. Since $Spin(8)$ is a Lie subgroup of F_4, the root system of $Spin(8)$ is contained in $\Phi(F_4)$, which is given as:

$$\Phi(Spin(8)) = \{\pm(t_i + t_j), \pm(t_i - t_j) \mid 1 \le i < j \le 4\}$$

It consists of the all long roots of the root system $\Phi(F_4)$. Then the Weyl group $W(Spin(8))$ is generated by the reflections associated with the long roots, and $W(Spin(8))$ is a subgroup of $W(F_4)$.

Put $W = W(Spin(8))$. The vertex set $W(F_4)$ of the GKM graph of F_4/T is decomposed into 6 cosets by the next theorem.

Theorem 3.1 ([A, Theorem 14.2])**.** *The Weyl group W of $Spin(8)$ is a normal subgroup of $W(F_4)$ and there is an isomorphism $W(F_4)/W \cong \mathfrak{S}_3$, where $\mathfrak{S}_n$ is the symmetric group on n-letters. Moreover $W(F_4)/W$ acts by permuting the three root pairs*

$$\pm\frac{1}{2}(t_1+t_2+t_3-t_4), \quad \pm\frac{1}{2}(t_1+t_2+t_3+t_4), \quad \pm t_4. \tag{3.1}$$

Let us describe the representatives of the cosets $W(F_4)/W$. First we define an element ρ of $W(F_4)$ as

$$\rho = \sigma_{\alpha_3}\sigma_{\alpha_2}\sigma_{\alpha_1}\sigma_{\alpha_0}\sigma_{\alpha_3}\sigma_{\alpha_2}\sigma_{\alpha_1}\sigma_{\alpha_3}\sigma_{\alpha_2}\sigma_{\alpha_4}, \tag{3.2}$$

where α_0 denotes the root $t_1 - t_2$ of $Spin(8)$. By a straightforward calculation, we have

$$\rho t_i = \begin{cases} -\gamma + t_i & (i = 1,2,3) \\ \gamma - t_4 & (i = 4), \end{cases} \tag{3.3}$$

$$\rho^2 t_i = \begin{cases} -\gamma + t_4 + t_i & (i = 1,2,3) \\ -\gamma & (i = 4), \end{cases}$$

and

$$\rho^3 = \mathrm{id}.$$

By the above equations the root system $\Phi(F_4)$ can be rewritten as:

$$\Phi(F_4) = \{\pm(t_i+t_j), \pm(t_i-t_j), \pm\rho^{\varepsilon}t_k \mid 1 \le i < j \le 4,\ 1 \le k \le 4,\ 0 \le \varepsilon \le 2\}$$

Note that ρ permutes the three root pairs (3.1) cyclically and $\kappa = \sigma_{t_4}$ interchanges $\pm\frac{1}{2}(t_1+t_2+t_3-t_4) = \pm\rho t_4$ and $\pm\frac{1}{2}(t_1+t_2+t_3+t_4) = \pm\rho^2 t_4$. Hence $W(F_4)/W \cong \mathfrak{S}_3$ is generated by ρ and κ. Since the equation

$$\kappa\rho = \rho^2\kappa \tag{3.4}$$

holds, we have a coset decomposition

$$W(F_4) = \coprod_{\substack{\varepsilon=0,1,2 \\ \delta=0,1}} \rho^{\varepsilon}\kappa^{\delta}W.$$

We will describe the GKM graph $\mathcal{F}_4$ of F_4/T. There are $24(=\#\Phi(F_4))$ edges out of each vertex of $\mathcal{F}_4$. The half of these edges correspond to the long roots $\pm(t_i \pm t_j)$ and the other half correspond to the short roots $\pm\rho^{\varepsilon}t_i$.

The subgraph induced by W is the GKM graph $\mathcal{G}$ of $Spin(8)/T$ and it is well understood in [FIM]. Let $\rho^{\varepsilon}\kappa^{\delta}\mathcal{G}$ be the GKM subgraph induced by $\rho^{\varepsilon}\kappa^{\delta}W$ for $\varepsilon = 0,1,2$ and $\delta = 0,1$. For any ε and δ the induced subgraph $\rho^{\varepsilon}\kappa^{\delta}\mathcal{G}$ is isomorphic to $\mathcal{G}$ as graphs. Indeed if an edge ww' in $\mathcal{G}$

satisfies $w' = w\sigma_\alpha$ for a root α of $Spin(8)$, then $\rho^\varepsilon\kappa^\delta w$ and $\rho^\varepsilon\kappa^\delta w'$ satisfy $\rho^\varepsilon\kappa^\delta w' = \rho^\varepsilon\kappa^\delta w\sigma_\alpha$, and vice versa. Moreover, labels of edges of $\rho^\varepsilon\kappa^\delta\mathcal{G}$ are also determined by $\mathcal{G}$ as follows. When an edge ww' has a root $\pm\beta$ as its label, the label of the edge connecting $\rho^\varepsilon\kappa^\delta w$ and $\rho^\varepsilon\kappa^\delta w'$ is $\pm\rho^\varepsilon\kappa^\delta\beta$. Remark that if an edge ww' in $\rho^\varepsilon\kappa^\delta\mathcal{G}$ satisfies $w' = w\sigma_\alpha$, then α is one of the long roots.

From the above argument, it is sufficient to consider the edges connecting two of $\rho^\varepsilon\kappa^\delta\mathcal{G}$'s, which correspond to the short roots. Easy calculations show that

$$\sigma_{t_4} = \kappa, \quad \sigma_{\rho t_4} = \rho^2\kappa, \quad \sigma_{\rho^2 t_4} = \rho\kappa.$$

Then the GKM graph $\mathcal{F}_4$ has an induced subgraph below, where e denotes the unit element of $W(F_4)$ and element of $W(F_4)$ in each circle denotes a vertex of $\mathcal{F}_4$. The labels are calculated later.



(3.5)

We will calculate the reflection σ_α for a short root α to describe $\mathcal{F}_4$. For example let us consider the short root ρt_1 and the reflection $\sigma_{\rho t_1}$. By (3.3) we have $\rho t_1 = \frac{1}{2}(t_1 - t_2 - t_3 - t_4) = \sigma_{t_2}\sigma_{t_3}(\rho t_4)$. Then $\sigma_{\rho t_1} = \sigma_{t_2}\sigma_{t_3}\sigma_{\rho t_4}\sigma_{t_3}\sigma_{t_2}$ and $\sigma_{t_2}\sigma_{t_3} \in W$. Since W is a normal subgroup of $W(F_4)$, we have $W \cdot \rho^2\kappa W = \rho^2\kappa W$ in $W(F_4)/W$. Hence $\sigma_{\rho t_1}$ is also contained in $\rho^2\kappa W$. For any i, it is shown similarly that

$$\sigma_{\rho t_i} \in \rho^2\kappa W, \quad \sigma_{\rho^2 t_i} \in \rho\kappa W$$

and obviously we have

$$\sigma_{t_i} \in \kappa W.$$

Hence, for any $0 \leq \varepsilon, \varepsilon' \leq 2$ and $\delta = 0, 1$, it is independent from the choice of i and $w \in \rho^{\epsilon'}\kappa^\delta W$ which coset contains $w\sigma_{\rho^\varepsilon t_i}$. We have understood the underlying graph of $\mathcal{F}_4$.

Let us calculate the label of the edge connecting the vertices κ and ρ in the GKM subgraph (3.5), which corresponds a short root ρt_4. Then the label of the edge is $\pm\kappa(\rho t_4)$. It follows from the relation (3.4) that

$$\pm\kappa(\rho t_4) = \pm\rho^2\kappa t_4 = \pm\rho^2 t_4.$$

One can make similarly calculations of the labels of other edges in the GKM subgraph (3.5). For any $w \in W$, w fixes three sets of short roots $\{\pm t_i\}_{i=1}^4$, $\{\pm\rho t_i\}_{i=1}^4$ and $\{\pm\rho^2 t_i\}_{i=1}^4$ since

w permutes t_i's and changes the signs of even number of t_i's. Hence the label $\pm\rho^{\varepsilon}\kappa^{\delta}w(\alpha)$ is calculated similarly for any short root α.

We can now describe a schematic diagram of $\mathcal{F}_4$ as below.


$\mathcal{G}$
$(\rho t_i, \sigma_{\rho t_j})$
$(\rho^2 t_i, \sigma_{\rho^2 t_j})$
$\rho^2\kappa\mathcal{G}$
$\rho\kappa\mathcal{G}$
$(\rho^2 t_i, \sigma_{t_j})$
$(\rho t_i, \sigma_{t_j})$
$(t_i, \sigma_{\rho^2 t_j})$
$(t_i, \sigma_{\rho t_j})$
$\rho\mathcal{G}$
(t_i, σ_{t_j})
$\rho^2\mathcal{G}$
$(\rho^2 t_i, \sigma_{\rho t_j})$
$(\rho t_i, \sigma_{\rho^2 t_j})$
$\kappa\mathcal{G}$

(3.6)

This diagram means the followings. For example, $\mathcal{G}$ and $\rho\mathcal{G}$ are not adjacent in this diagram. It means that for any vertices $w \in W$ and $w' \in \rho W$, they are not adjacent. On the other hand, $\rho\mathcal{G}$ and $\rho\kappa\mathcal{G}$ are adjacent in this diagram, and a pair $(\rho t_i, \sigma_{t_j})$ is assigned to the edge. The first entry ρt_i is a root and the second entry σ_{t_j} is a reflection. If two vertices $w \in \rho W$ and $w' \in \rho\kappa W$ are adjacent in $\mathcal{F}_4$, then they satisfy $w' = w\sigma_{t_j}$ for some j, and the edge ww' is labeled by ρt_i for some i. The label $\pm\rho t_i$ equals to $\pm w t_j$. Especially each vertex of $\rho\mathcal{G}$ is connected to 4 vertices of $\rho\kappa\mathcal{G}$ by the edges correspond to the short roots t_j's $(1 \leq j \leq 4)$, and vice versa. The labels of these edges are $\pm\rho t_i$'s $(1 \leq i \leq 4)$. Every ρt_i's appear as the labels of the edges out of each vertex of $\rho\mathcal{G}$. The situation is the same for any connected two subgraphs in the schematic diagram (3.6).

4 Proof of the main theorem

There is a fibration sequence

$$F_4/T \longrightarrow ET \times_T F_4/T \longrightarrow BT. \tag{4.1}$$

The cohomology rings of F_4/T and BT are free as $\mathbb{Z}$-modules and have vanishing odd parts. As shown in Section 3, $H^*(BT)$ has 5 generators t_1, t_2, t_3, t_4, and γ of degree 2 and one relation of degree 2. According to [TW], $H^*(F_4/T)$ has τ_1, τ_2, τ_3, τ_4, and γ_1 of degree 2, γ_3 of degree 6, and ω of degree 8 as its generators, and $H^*(F_4/T)$ has seven relations of degree 2, 4, 6, 8, 12, 16, and 24. We can expect $H^*_T(F_4/T)$ has corresponding generators and relations. It is easy to see the Poincaré series of F_4/T and BT are

$$(1 + x^8 + x^{16}) \prod_{i=1}^{4} \frac{1 - x^{4i}}{1 - x^2} \quad \text{and} \quad \frac{1}{(1 - x^2)^4},$$

respectively. Hence we obtain the following proposition by Serre spectral sequence.

Proposition 4.1. *$H_T^*(F_4/T)$ is free as a $\mathbb{Z}$-module and its Poincaré series is*

$$P(H^*(ET\times_T F_4/T),x)=\frac{1}{(1-x^2)^4}(1+x^8+x^{16})\prod_{i=1}^{4}\frac{1-x^{4i}}{1-x^2}.$$

By the fibration sequence (4.1) generators of $H_T^*(F_4/T)$ come from the cohomology of F_4/T or BT. Let us define the corresponding GKM functions t_i, γ, τ_i, γ_1 and $\gamma_3\in\mathrm{Map}(W(F_4),H^*(BT))$ for $1\le i\le 4$ and GKM functions γ_2 and γ_4 for convenience in Section 5 as: for any $w\in W(F_4)$

$$\begin{aligned}t_i(w)&=t_i\ \ (i=1,\dots,4)\\ \gamma(w)&=\gamma\\ \tau_i(w)&=w(t_i)\ \ (i=1,\dots,4)\\ \gamma_j&=\frac{1}{2}(e_j(\tau)-e_j(t))\ \ (j=1,2,3),\end{aligned}$$

and

$$\gamma_4(w)=\begin{cases}0 & w\in W\sqcup\rho^2\kappa W\\ e_4(\rho^2t) & w\in\rho^2W\sqcup\rho\kappa W\\ -e_4(t) & w\in\rho W\sqcup\kappa W.\end{cases}\tag{4.2}$$

Moreover we define $\omega=e_4(\tau)-e_4(t)-2\gamma_4$. Then

$$\omega(w)=\begin{cases}0 & w\in W\sqcup\kappa W\\ -e_4(\rho^2t) & w\in\rho W\sqcup\rho\kappa W\\ e_4(\rho t) & w\in\rho^2W\sqcup\rho^2\kappa W\end{cases}\tag{4.3}$$

Since t_i's and γ are constant functions, they are GKM functions. A straightforward calculation shows the following relation holds.

$$e_4(t)+e_4(\rho t)+e_4(\rho^2t)=0\tag{4.4}$$

By the schematic diagram (3.6) of $\mathcal{F}_4$ one can see γ_4 is a GKM function since $e_4(\rho^\varepsilon t)$ is the product of all $\rho^\varepsilon t_1$, $\rho^\varepsilon t_2$, $\rho^\varepsilon t_3$, and $\rho^\varepsilon t_4$ for $\varepsilon=0,1,2$. The following calculation shows τ_i's satisfy the GKM condition. For any edge ww' which satisfies $w'=w\sigma_\alpha$, we have

$$\begin{aligned}\tau_i(w)-\tau_i(w')&=w(t_i)-w'(t_i)\\ &=w\Big(t_i-\Big(t_i-2\frac{(t_i,\alpha)}{(\alpha,\alpha)}\alpha\Big)\Big)\\ &=2\frac{(t_i,\alpha)}{(\alpha,\alpha)}w\alpha.\end{aligned}$$

Since GKM functions form a ring, for $j=1,2,3$, we see that γ_j's are functions from $W(F_4)$ to $H^*(BT)\otimes\mathbb{Z}[\frac{1}{2}]$ which satisfy the GKM condition tensoring with $\mathbb{Z}[\frac{1}{2}]$. The following calculations show that γ_j's are $H^*(BT)$-valued functions. Let ρ extend to an automorphism of $H^*(BT)$ naturally. For $w \in W \sqcup \kappa W = W(Spin(9))$ and $\varepsilon = 0, 1, 2$,

$$\begin{aligned}
\gamma_j(\rho^\varepsilon w) &= \frac{1}{2}(e_j(\tau) - e_j(t))(\rho^\varepsilon w) \\
&= \frac{1}{2}(\rho^\varepsilon e_j(w(t)) - e_j(t)) \\
&= \rho^\varepsilon\Big(\frac{1}{2}(e_j(w(t)) - e_j(t))\Big) + \frac{1}{2}(e_j(\rho^\varepsilon t) - e_j(t)).
\end{aligned}$$

Since w only permutes t_i's and changes their signs, it is obvious that $\frac{1}{2}(e_j(w(t)) - e_j(t)) \in H^*(BT)$. Then $\rho^\varepsilon(\frac{1}{2}(e_j(w(t)) - e_j(t))) \in H^*(BT)$. On the other hand one can see $\frac{1}{2}(e_j(\rho^\varepsilon t) - e_j(t)) \in H^*(BT)$ for $\varepsilon = 0, 1, 2$ as follows. When $\varepsilon = 0$, $\frac{1}{2}(e_j(\rho^\varepsilon t) - e_j(t)) = 0$ and it is contained in $H^*(BT)$. When $\varepsilon = 1, 2$, Table 1 shows the value of $\frac{1}{2}(e_j(\rho^\varepsilon t) - e_j(t))$ for $j = 1, 2, 3$. Then γ_j is a $H^*(BT)$-valued function and then a GKM function. The following lemma will be proved in Section 5.

Lemma 4.1 (see [FIM, Lemma 5.4]). *Let $\mathcal{F}_4$ be the GKM graph of F_4/T, then $H^*(\mathcal{F}_4)$ is generated by the GKM functions t_i, γ, τ_i, γ_1, γ_3, ω $(i = 1, 2, 3, 4)$ as a ring.*

Table 1: the value of $\frac{1}{2}(e_j(\rho^\varepsilon t) - e_j(t))$

	$j = 1$	$j = 2$	$j = 3$
$\varepsilon = 1$	$-\gamma - t_4$	$-\gamma^2 + t_4$	$t_4\gamma(\gamma - t_4) - t_4(t_1t_2 + t_2t_3 + t_3t_1)$
$\varepsilon = 2$	$-2\gamma + t_4$	$(-2\gamma + t_4)t_4$	$\gamma^3 - t_4\gamma^2 - \gamma(t_1t_2 + t_2t_3 + t_3t_1)$

By the fibration sequence (4.1) we can expect some relations hold in $H^*(\mathcal{F}_4)$, which come from the relations of $H^*(BT)$ and $H^*(F_4/T)$. Proposition 4.2 claims the corresponding relations hold in $H^*(\mathcal{F}_4)$.

Proposition 4.2. *The following relations hold in $H^*(\mathcal{F}_4) \subset \mathrm{Map}(W(F_4), H^*(BT))$;*

$$2\gamma = e_1(t) \tag{4.5}$$

$$2\gamma_1 = e_1(\tau) - e_1(t) \tag{4.6}$$

$$2\gamma_1^2 - (e_2(\tau) - e_2(t)) + (e_1(\tau) - e_1(t))e_1(t) \tag{4.7}$$

$$2\gamma_3 = e_3(\tau) - e_3(t) \tag{4.8}$$

$$2A + e_4(\tau) - e_4(t) - 3\omega = 0 \tag{4.9}$$

$$\gamma_3^2 + \gamma_3 e_3(t) - \gamma_1^2 e_4(\tau) - \gamma_1 e_1(t) e_4(\tau) - \gamma_4 e_2(t) - \gamma^2\omega = 0 \tag{4.10}$$

$$e_4(\tau)^2 - A^2 + (e_4(\tau) + 2\omega)A - e_4(\tau)(2\omega + e_4(t)) - \omega e_4(\rho t) = 0 \tag{4.11}$$

$$\omega(\omega - e_4(\rho t))(\omega + e_4(\rho^2 t)) = 0, \tag{4.12}$$

where $A = \gamma_1^4 + \gamma_1^2 e_1(t)e_1(\tau) - \gamma_3 e_1(\tau) - \gamma_1 e_3(t) + \gamma_1^2 e_2(t) + \gamma_1 e_1(t) e_2(t)$.

We have also the following relations in $H^*(\mathcal{F}_4)$;

$$2\gamma_2 = e_2(\tau) - e_2(t) \tag{4.13}$$
$$\gamma_2 = \gamma_1^2 + \gamma_1 e_1(t) \tag{4.14}$$
$$\gamma_2^2 - \gamma_3 e_1(\tau) - \gamma_1 e_3(t) + \gamma_2 e_2(t) - e_4(\tau) + e_4(t) + 3\gamma_4 = 0 \tag{4.15}$$
$$\gamma_4(\gamma_4 + e_4(t))(\gamma_4 - e_4(\rho^2 t)) = 0 \tag{4.16}$$

Proposition 4.2 and these relations (4.13), (4.14), (4.15), (4.16) are proved in Section 6. The equation (4.15) can be modified as:

$$\begin{aligned} \gamma_4 &= \omega - \gamma_2^2 + \gamma_3 e_1(\tau) + \gamma_1 e_3(t) - \gamma_2 e_2(t) \\ &= \omega - A \end{aligned} \tag{4.17}$$

This relation claims that γ_4 can replace ω as a generator of $H^*(\mathcal{F}_4)$. The following lemma is proved in Section 7.

Lemma 4.2. *$\mathbb{Z}[t_i, \gamma, \tau_i, \gamma_1, \gamma_3, \omega]/(r_1', r_1, r_2, r_3, r_4, r_6, r_8, r_{12})$ is free as a $\mathbb{Z}$-module, and its Poincaré series coincides with that of $H_T^*(F_4/T)$.*

Now we are ready to prove Theorem 1.1.

Proof of Theorem 1.1. Let I denote the ideal $(r_1', r_1, r_2, r_3, r_4, r_6, r_8, r_{12})$ in the polynomial ring $\mathbb{Z}[t_i, \gamma, \tau_i, \gamma_1, \gamma_3, \omega \mid 1 \le i \le 4]$. We have a surjective ring homomorphism

$$\mathbb{Z}[t_i, \gamma, \tau_i, \gamma_1, \gamma_3, \omega \mid 1 \le i \le 4] \to H^*(\mathcal{F}_4)$$

by Lemma 4.1, and it factors through $\mathbb{Z}[t_i, \gamma, \tau_i, \gamma_1, \gamma_3, \omega \mid 1 \le i \le 4]/I \to H^*(\mathcal{F}_4)$ by Proposition 4.2. It follows from Proposition 4.1 and Lemma 4.2 that $H^*(\mathcal{F}_4)$ and $\mathbb{Z}[t_i, \gamma, \tau_i, \gamma_1, \gamma_3, \omega \mid 1 \le i \le 4]/I$ are free as $\mathbb{Z}$-modules, and $\mathbb{Z}[t_i, \gamma, \tau_i, \gamma_1, \gamma_3, \omega \mid 1 \le i \le 4]/I$ and $H^*(\mathcal{F}_4)$ have the same rank in each degree. Therefore the ring homomorphism $\mathbb{Z}[t_i, \gamma, \tau_i, \gamma_1, \gamma_3, \omega \mid 1 \le i \le 4]/I \to H^*(\mathcal{F}_4)$ is an isomorphism and Theorem 1.1 is proved by Theorem 2.1. $\square$

5 Proof of Lemma 4.1

First we introduce some notation for the proof of Lemma 4.1. For $n \in \mathbb{N}$ let $[n]$ and $\pm[n]$ be $\{i \in \mathbb{Z} \mid 1 \le i \le n\}$ and $\{\pm i \in \mathbb{Z} \mid 1 \le i \le n\}$, respectively. For $1 \le n \le 4$, let I_n denote an ordered n-tuple $(i_1, \dots, i_n)$ of elements of $[4]$ which does not include the same entries, and I_n' denote an ordered n-tuple $(i_1', \dots, i_n')$ of elements of $\pm[4]$ such that $|i_k'| \ne |i_l'|$ for $k \ne l$. We often regard I_n, I_n' as the non-ordered n-subsets of $[4]$ by the following maps.

$$(i_1, \dots, i_n) \mapsto \{i_1, \dots, i_n\}, \quad (i_1', \dots, i_n') \mapsto \{|i_1'|, \dots, |i_n'|\}$$

For $\varepsilon = 0, 1, 2$, we define a subset $\rho^\varepsilon W_{I_n'}^{I_n}$ of $W(F_4)$ as:

$$\rho^\varepsilon W_{I_n'}^{I_n} = \{w \in W(F_4) \mid w \in \rho^\varepsilon W(Spin(9)),\ w(t_{i_k}) = \rho^\varepsilon t_{i_k'}\ (1 \le k \le n)\}$$

We define I_0 and I'_0 to be the empty set. Note that $\rho^\varepsilon W^{I_{n-1}}_{I'_{n-1}}$ includes $\rho^\varepsilon W^{I_n}_{I'_n}$ and decomposes as follows.

$$\rho^\varepsilon W^{I_{n-1}}_{I'_{n-1}} = \coprod_{i_n \in [4]\setminus I_{n-1}} \rho^\varepsilon W^{(I_{n-1},i_n)}_{(I'_{n-1},i'_n)} \sqcup \coprod_{i_n \in [4]\setminus I_{n-1}} \rho^\varepsilon W^{(I_{n-1},i_n)}_{(I'_{n-1},-i'_n)} \tag{5.1}$$

For a set $S = \{j_1, \ldots, j_k\}$ of natural numbers with $j_1 < \cdots < j_k$, let x_S denote a sequence $(x_{j_1}, \ldots, x_{j_k})$ for $x = t,\ \rho t,\ \rho^2 t,\ \tau$. For $n \geq 0$, $j \leq 4$, and $\varepsilon = 0, 1, 2$, let $\gamma^{(\varepsilon)}_j{}^{I_n}_{I'_n}$ be a function from $\rho^\varepsilon W^{I_n}_{I'_n}$ to $\mathbb{Z}[\frac{1}{2}][t_1, t_2, t_3, t_4]$ defined as

$$\gamma^{(\varepsilon)}_j{}^{I_n}_{I'_n} = \frac{1}{2}(e_j(\tau_{[4]\setminus I_n}) - e_j(\rho^\varepsilon t_{[4]\setminus I'_n})),$$

where I_n and I'_n are regarded as subsets of [4]. When $n = 0$ we abbreviate $\gamma^{(\varepsilon)}_j{}^{\emptyset}_{\emptyset}$ by $\gamma^{(\varepsilon)}_j$. If $j \leq 0$ or $j > 4 - n$, we define $\gamma^{(\varepsilon)}_j{}^{I_n}_{I'_n} = 0$.

By the relation (4.17) and (4.15), it is sufficient for proving Lemma 4.1 to show that GKM functions t_i, γ, τ_i, γ_i $(1 \leq i \leq 4)$ generate $H^*(\mathcal{F}_4)$ as a ring. Let t_{-i} denote $-t_i$. We define a function $f^{(\varepsilon)}{}^{I_{n-1}}_{i'_n}$ which is useful in the proof of Lemma 4.1 as:

$$f^{(\varepsilon)}{}^{I_{n-1}}_{i'_n} = \frac{1}{2} \prod_{k \in [4]\setminus I_{n-1}} (\tau_k - \rho^\varepsilon t_{i'_n}).$$

This function is $H^*(BT)$-valued on $\rho^\varepsilon W^{I_{n-1}}_{I'_{n-1}}$, since for any $w \in \rho^\varepsilon W^{I_{n-1}}_{I'_{n-1}}$ there exists $k \in [4] \setminus I_{n-1}$ such that $w \in \rho^\varepsilon W^{(I_{n-1},k)}_{(I'_{n-1},i'_n)} \sqcup \rho^\varepsilon W^{(I_{n-1},k)}_{(I'_{n-1},-i'_n)}$ by the decomposition (5.1), and then $w(t_k) - \rho^\varepsilon t_{i'_n}$ equals to 0 or $-2\rho^\varepsilon t_{i'_n}$. Especially this function vanishes on the summand $\coprod_{k\in[4]\setminus I_{n-1}} \rho^\varepsilon W^{(I_{n-1},k)}_{(I'_{n-1},i'_n)}$ and for $w \in \rho^\varepsilon W^{(I_{n-1},k)}_{(I'_{n-1},-i'_n)}$

$$f^{(\varepsilon)}{}^{I_{n-1}}_{i'_n}(w) = -\rho^\varepsilon t_{i'_n} \prod_{k \in [4]\setminus I'_n} (\rho^\varepsilon t_k - \rho^\varepsilon t_{i'_n}). \tag{5.2}$$

Let R denote the subring of $H^*(\mathcal{F}_4)$ generated by t_i's, γ, τ_i's, and γ_i's $(1 \leq i \leq 4)$. The following proposition claims this function $f^{(\varepsilon)}{}^{I_{n-1}}_{i'_n}$ can be replaced partly by an element of R.

Proposition 5.1. *For $1 \leq n \leq 4$, there is a polynomial in γ_1, γ_2, γ_3, γ_4 over $H^*(BT)$, which coincides with the function $f^{(\varepsilon)}{}^{I_{n-1}}_{i'_n}$ on $\rho^\varepsilon W^{I_{n-1}}_{I'_{n-1}}$.*

Proposition 5.1 is a consequence by Lemma 5.1 and 5.2 below.

Lemma 5.1. *For $1 \leq n \leq 4$, there is a polynomial in $\gamma^{(\varepsilon)}_j{}^{I_{n-1}}_{I'_{n-1}}$'s $(1 \leq j \leq 4 - (n-1))$ over $H^*(BT)$, which coincides with $f^{(\varepsilon)}{}^{I_{n-1}}_{i'_n}$ on $\rho^\varepsilon W^{I_{n-1}}_{I'_{n-1}}$.*

Lemma 5.2 (see [FIM, Lemma 5.3]). *For $1 \leq n \leq 4$, there is a polynomial in $\gamma^{(\varepsilon)I_{n-1}}_{1\ \,I'_{n-1}}, \ldots, \gamma^{(\varepsilon)\ \,I_{n-1}}_{4-n\,I'_{n-1}}$ over $H^*(BT)$, which coincides with $\gamma^{(\varepsilon)(I_{n-1},i_n)}_{j\ \,(I'_{n-1},i'_n)}$ on $\rho^\varepsilon W^{(I_{n-1},i_n)}_{(I'_{n-1},i'_n)}$. More explicitly,*

$$\gamma^{(\varepsilon)I_n}_{j\ \,I'_n} = \begin{cases} \sum_{k=0}^{j-1} \gamma^{(\varepsilon)\ \,I_{n-1}}_{j-k\,I'_{n-1}}(-\rho^\varepsilon t_{i'_n})^k & \operatorname{sgn} i'_n = 1 \\ \sum_{k=0}^{j-1} \gamma^{(\varepsilon)\ \,I_{n-1}}_{j-k\,I'_{n-1}}(-\rho^\varepsilon t_{i'_n})^k + \sum_{k=1}^{j} e_{j-k}(\rho^\varepsilon t_{[4]\setminus I'_n})(-\rho^\varepsilon t_{i'_n})^k & \operatorname{sgn} i'_n = -1. \end{cases}$$

Proof of Proposition 5.1. By Lemma 5.1, there is a polynomial in $\gamma^{(\varepsilon)I_{n-1}}_{j\ \,I'_{n-1}}$'s $(1 \leq j \leq 4-(n-1))$ over $H^*(BT)$, which coincides with $f^{(\varepsilon)I_{n-1}}_{\ \ i'_n}$ on $\rho^\varepsilon W^{I_{n-1}}_{I'_{n-1}}$ for $\varepsilon = 0, 1, 2$. Then by Lemma 5.2 $\gamma^{(\varepsilon)(I_{n-1},i_n)}_{j\ \,(I'_{n-1},i'_n)}$ can be replaced by some polynomial in $\gamma^{(\varepsilon)I_{n-1}}_{1\ \,I'_{n-1}}, \ldots, \gamma^{(\varepsilon)\ \,I_{n-1}}_{4-n\,I'_{n-1}}$ over $H^*(BT)$. Iterated use of Lemma 5.2 makes n smaller and smaller, and finally we reached to a polynomial in γ^ε_1, γ^ε_2, γ^ε_3, γ^ε_4 over $H^*(BT)$, which coincides with $f^{(\varepsilon)I_{n-1}}_{\ \ i'_n}$ on $\rho^\varepsilon W^{I_{n-1}}_{I'_{n-1}}$ for $\varepsilon = 0, 1, 2$. Next we need to show that $\gamma_j - \gamma^{(\varepsilon)}_j \in H^*(BT)$ on $\rho^\varepsilon W(Spin(9))$ for $1 \leq j \leq 4$ and $\varepsilon = 0, 1, 2$ to complete the proof of Proposition 5.1. By definition we have

$$\gamma^{(\varepsilon)}_j = \gamma_j + \frac{1}{2}(e_j(t) - e_j(\rho^\varepsilon t)) \quad (j = 1, 2, 3). \tag{5.3}$$

For $\varepsilon = 0, 1, 2$ and $j = 1, 2, 3$, Table 1 shows that $(e_j(t) - e_j(\rho^\varepsilon t))/2 \in H^*(BT)$ and then $\gamma_j - \gamma^{(0)}_{\varepsilon,j} \in H^*(BT)$ on $\rho^\varepsilon W(Spin(9))$. By the definition of γ_4 and the equation (4.4), we have

$$\begin{aligned} \gamma^{(0)}_4 &= \gamma_4 && \text{on } W(Spin(9)), \\ \gamma^{(1)}_4 &= \gamma_4 + e_4(t) && \text{on } \rho W(Spin(9)), \\ \gamma^{(2)}_4 &= \gamma_4 - e_4(\rho^2 t) && \text{on } \rho^2 W(Spin(9)). \end{aligned}$$

Therefore there is a polynomial in γ_1, γ_2, γ_3, γ_4 over $H^*(BT)$, which coincides with the function $f^{(\varepsilon)I_{n-1}}_{\ \ i'_n}$ on $\rho^\varepsilon W^{I_{n-1}}_{I'_{n-1}}$. □

Proof of Lemma 5.1. Without loss of generality, we may suppose that $I_{n-1} = (1, \ldots, n-1)$. Note that $e_j(x_S) = 0$ for $j > \#S$ or $j < 0$ and we have

$$e_j(x_1, \ldots, x_{m-1}, x_m) = e_j(x_1, \ldots, x_{m-1}) + e_{j-1}(x_1, \ldots, x_{m-1})x_m. \tag{5.4}$$

By the definition of $\gamma^{(\varepsilon)I_{n-1}}_{j\ \,I'_{n-1}}$ we can modify the GKM function

$$\frac{1}{2}\prod_{l=0}^{4-n}(\tau_{n+l} - \rho^\varepsilon t_{i'_n}) = \frac{1}{2}\sum_{j=0}^{5-n} e_j(\tau_{[4]\setminus I_{n-1}})(-\rho^\varepsilon t_{i'_n})^{5-n-j}$$

to

$$\frac{1}{2}\sum_{j=0}^{5-n}(2\gamma^{(\varepsilon)I_{n-1}}_{j\ \,I'_{n-1}} + e_j(\rho^\varepsilon t_{[4]\setminus I'_{n-1}}))(-\rho^\varepsilon t_{i'_n})^{5-n-j}. \tag{5.5}$$

Pay attention to the sign of i'_n and recall $[4] \setminus I'_{n-1} = \{i \in [4] \mid \pm i \notin I'_{n-1}\}$. By the equation (5.4), (5.5) equals to

$$\begin{aligned}
&\sum_{j=0}^{5-n} \gamma_{j\;\;I'_{n-1}}^{(\varepsilon)\,I_{n-1}} (-\rho^\varepsilon t_{i'_n})^{5-n-j} + \frac{1}{2}\sum_{j=0}^{5-n} (e_j(\rho^\varepsilon t_{[4]\setminus I'_n}) + e_{j-1}(\rho^\varepsilon t_{[4]\setminus I'_n})\rho^\varepsilon t_{|i'_n|})(-\rho^\varepsilon t_{i'_n})^{5-n-j} \\
&\quad = \begin{cases} \sum\limits_{j=0}^{5-n} \gamma_{j\;\;I'_{n-1}}^{(\varepsilon)\,I_{n-1}} (-\rho^\varepsilon t_{i'_n})^{5-n-j} & \operatorname{sgn} i'_n = 1 \\ \sum\limits_{j=0}^{5-n} \gamma_{j\;\;I'_{n-1}}^{(\varepsilon)\,I_{n-1}} (-\rho^\varepsilon t_{i'_n})^{5-n-j} + \sum\limits_{j=0}^{4-n} e_j(\rho^\varepsilon t_{[4]\setminus I'_n})(-\rho^\varepsilon t_{i'_n})^{5-n-j} & \operatorname{sgn} i'_n = -1. \end{cases}
\end{aligned}$$

This calculation shows Lemma 5.1. ☐

Proof of Lemma 5.2. The relation $\tau_{i_n} = \rho^\varepsilon t_{i'_n}$ holds on $\rho^\varepsilon W^{(I_{n-1}, i_n)}_{(I'_{n-1}, i'_n)}$. Then we have

$$\begin{aligned}
&\gamma_{j\;\;I'_{n-1}}^{(\varepsilon)\,I_{n-1}} - \gamma_{j\;\;(I'_{n-1}, i'_n)}^{(\varepsilon)\,(I_{n-1}, i_n)} \\
&= \frac{1}{2}(e_j(\tau_{i \in [4]\setminus I_{n-1}}) - e_j(\rho^\varepsilon t_{i' \in [4]\setminus I'_{n-1}})) - \frac{1}{2}(e_j(\tau_{i \in [4]\setminus I_n}) - e_j(\rho^\varepsilon t_{i' \in [4]\setminus I'_n})) \\
&= \frac{1}{2}(e_{j-1}(\tau_{i \in [4]\setminus I_n})\tau_{i_n} - e_{j-1}(\rho^\varepsilon t_{i' \in [4]\setminus I'_n})\rho^\varepsilon t_{|i'_n|}) \\
&= \begin{cases} \gamma_{j-1\;I'_n}^{(\varepsilon)\;\;I_n} \rho^\varepsilon t_{i'_n} & \operatorname{sgn} i'_n = 1 \\ \gamma_{j-1\;I'_n}^{(\varepsilon)\;\;I_n} \rho^\varepsilon t_{i'_n} + e_{j-1}(\rho^\varepsilon t_{i' \in [4]\setminus I'_n})\rho^\varepsilon t_{i'_n} & \operatorname{sgn} i'_n = -1. \end{cases}
\end{aligned}$$

Iterated use of this equation shows that

$$\gamma_{j\;\;(I'_{n-1}, i'_n)}^{(\varepsilon)\,(I_{n-1}, i_n)} = \begin{cases} \sum\limits_{k=0}^{j-1} \gamma_{j-k\;I'_{n-1}}^{(\varepsilon)\;\;I_{n-1}} (-\rho^\varepsilon t_{i'_n})^k & \operatorname{sgn} i'_n = 1 \\ \sum\limits_{k=0}^{j-1} \gamma_{j-k\;I'_{n-1}}^{(\varepsilon)\;\;I_{n-1}} (-\rho^\varepsilon t_{i'_n})^k + \sum\limits_{k=1}^{j} e_{j-k}(\rho^\varepsilon t_{[4]\setminus I'_n})(-\rho^\varepsilon t_{i'_n})^k & \operatorname{sgn} i'_n = -1. \end{cases}$$

☐

Now we are ready to prove Lemma 4.1.

Proof of Lemma 4.1. We only need to show that any GKM function $h \in H^*(\mathcal{F}_4)$ belongs to the subring R generated by t_i's, γ, τ_i's, and γ_i's $(1 \leq i \leq 4)$. By the definition of ρ, the set of all vertices $W(F_4)$ of $\mathcal{F}_4$ decomposes as:

$$W(F_4) = W(Spin(9)) \sqcup \rho W(Spin(9)) \sqcup \rho^2 W(Spin(9)).$$

For each $\varepsilon = 0, 1, 2$, $\rho^\varepsilon W(Spin(9))$ has a filtration

$$\rho^\varepsilon W^{I_4}_{I'_4} \subset \cdots \subset \rho^\varepsilon W^{I_n}_{I'_n} \subset \rho^\varepsilon W^{I_{n-1}}_{I'_{n-1}} \subset \cdots \subset \rho^\varepsilon W^{I_0}_{I'_0} = \rho^\varepsilon W(Spin(9)).$$

By decreasing induction on n, we will show that any GKM function h can be modified to be 0 on $\rho^{\varepsilon} W^{I_n}_{I'_n}$ by subtracting some GKM function in R. Moreover in the induction step on n we give an induction to fill the decomposition (5.1) of $\rho^{\varepsilon} W^{I_{n-1}}_{I'_{n-1}}$.

Let $0 \leq n \leq 4$. The following claim in the case where $n = 0$ shows that h can be modified to be 0 on $W(Spin(9))$.

Claim 1 (n)**.** *For any ordered n-tuples I_n, I'_n and any function h from $W^{I_n}_{I'_n}$ to $H^*(BT)$ which satisfies the GKM condition on $W^{I_n}_{I'_n}$, there is a GKM function $G \in R$ which coincides with h on $W^{I_n}_{I'_n}$.*

We show this claim by decreasing induction on n. For $n = 4$, since $W^{I_4}_{I'_4}$ is a one point set, the claim holds obviously. Assume Claim 1 (n) holds, and fix $I_n = (i_1, \dots, i_n)$ and $I'_n = (i'_1, \dots, i'_n)$. Then subtracting the polynomial of Claim 1 (n) from h, which coincides with h on $W^{I_n}_{I'_n}$, we may assume h vanishes on $W^{I_n}_{I'_n}$. We give an induction to fill the decomposition (5.1) of $W^{I_{n-1}}_{I'_{n-1}}$ as follows. For any $k \in [4] \setminus I_n$ and $w \in W(k, i'_n)$, $w\sigma \in W(i_n, i'_n)$, where σ denotes the reflection associated with $t_k - t_{i'_n}$. Then by the GKM condition, $h(w) - h(w\sigma) = h(w)$ belongs to the ideal generated by $w(t_{i_n} - t_k) = w(t_{i_n}) - t_{i'_n} = \tau_{i_n}(w) - t_{i'_n}$. Put $k_0, \dots, k_{4-n} \in [4] \setminus I_{n-1}$ as $k_0 = i_n$, $k_s < k_t$ for $1 \leq s < t$ and $\{k_0, \dots, k_{4-n}\} \cup I_{n-1} = [4]$.

Claim 2 (t)**.** *If h is a GKM function which vanishes on $\coprod_{s<t} W^{(I_{n-1},k_s)}_{(I'_{n-1},i'_n)}$, there is a GKM function $G \in R$ such that h coincides with $\prod_{s<t}(\tau_{k_s} - t_{i'_n})G$ on $W^{(I_{n-1},k_t)}_{(I'_{n-1},i'_n)}$.*

We show this claim by induction on t $(0 \leq t \leq 4 - n)$. Without loss of generality, we may suppose that $I_{n-1} = (1, \dots, n-1)$ and $k_0 = i_n = n$, $k_1 = n+1, \dots, k_{4-n} = 4$. We rephrase Claim 2 (t) as:

Claim 2 (k)**.** *If h vanishes on $\coprod_{0 \leq l < k} W^{(I_{n-1},n+l)}_{(I'_{n-1},i'_n)}$, there is a GKM function $G \in R$ such that h coincides with $\prod_{0 \leq l < k}(\tau_{n+l} - t_{i'_n})G$ on $W^{(I_{n-1},n+k)}_{(I'_{n-1},i'_n)}$.*

Obviously $\prod_{0 \leq l < k}(\tau_{n+l} - t_{i'_n})$ vanishes on $\coprod_{0 \leq l < k} W^{(I_{n-1},n+l)}_{(I'_{n-1},i'_n)}$. For $w \in W^{(I_{n-1},n+k)}_{(I'_{n-1},i'_n)}$, by the GKM condition, there is an element $g_w \in H^*(BT)$ such that

$$h(w) = \prod_{0 \leq l < k} (\tau_{n+l} - t_{i'_n})(w) g_w.$$

One can verify that a function $G' \colon W^{(I_{n-1},n+k)}_{(I'_{n-1},i'_n)} \to H^*(BT)$ given by

$$G'(w) = g_w$$

satisfies the GKM condition on $W^{(I_{n-1},n+k)}_{(I'_{n-1},i'_n)}$ as follows. Assume that two vertices w, $w' \in W^{(I_{n-1},n+k)}_{(I'_{n-1},i'_n)}$ of $\mathcal{F}_4$ satisfy $w' = w\sigma_\alpha$ for some positive root α. Then $\alpha = t_i - t_j$, where $i < j$ and

i, $j \in \{m \in \mathbb{Z} \mid n \leq m \leq n+k-1 \text{ or } n+k+1 \leq m \leq 4\}$. When $i < j < n+k$ or $n+k < i < j$, the GKM condition says

$$\begin{aligned} h(w) - h(w') &= \prod_{0 \leq l < k} (w(t_{n+l}) - t_{i'_n}) G'(w) - \prod_{0 \leq l < k} (w\sigma_{t_i - t_j}(t_{n+l}) - t_{i'_n}) G'(w') \\ &= \prod_{0 \leq l < k} (w(t_{n+l}) - t_{i'_n})(G'(w) - G'(w')) \end{aligned}$$

belongs to the ideal $(w(t_i - t_j))$. Since $w(t_{n+l}) - t_{i'_n}$ and $w(t_i - t_j)$ are prime, $G'(w) - G'(w')$ also belongs to the ideal $(w(t_i - t_j))$. When $i < n+k < j$, the GKM condition says

$$\begin{aligned} & h(w) - h(w') \\ = & \prod_{0 \leq l < k} (w(t_{n+l}) - t_{i'_n}) G'(w) - \prod_{0 \leq l < k} (w\sigma_{t_i - t_j}(t_{n+l}) - t_{i'_n}) G'(w') \\ = & \prod_{0 \leq l < k,\ l \neq i} (w(t_{n+l}) - t_{i'_n}) \Big((w(t_i) - t_{i'_n})(G'(w) - G'(w')) + (w(t_i) - w(t_j)) G'(w') \Big) \end{aligned}$$

belongs to the ideal $(w(t_i - t_j))$. Since $w(t_{n+l}) - t_{i'_n}$ and $w(t_i - t_j)$ are prime, $G'(w) - G'(w')$ also belongs to the ideal $(w(t_i - t_j))$. Hence the function G' satisfies the GKM condition on $W^{(I_{n-1}, n+k)}_{(I'_{n-1}, i'_n)}$.

By (decreasing) induction on n there is a GKM function $G \in R$ such that G and G' coincide on $W^{(I_{n-1}, n+k)}_{(I'_{n-1}, i'_n)}$. Then

$$h - \prod_{0 \leq l < k} (\tau_{n+l} - t_{i'_n}) G = 0 \quad \text{on} \coprod_{0 \leq l \leq k} W^{(I_{n-1}, n+l)}_{(I'_{n-1}, i'_n)}.$$

Therefore the induction on k proceeds.

Next we fill the other half of the decomposition (5.1). Note that when $I_{n-1} = (1, \ldots, n-1)$,

$$f^{(0)}{}^{I_{n-1}}_{i'_n} = \frac{1}{2} \prod_{0 \leq l \leq 4-n} (\tau_{n+l} - t_{i'_n}).$$

Let $0 \leq k' \leq 4 - n$.

Claim 3 (k')**.** *If h vanishes on $\coprod_{0 \leq l \leq 4-n} W^{(I_{n-1}, n+l)}_{(I'_{n-1}, i'_n)} \sqcup \coprod_{0 \leq l < k'} W^{(I_{n-1}, n+l)}_{(I'_{n-1}, -i'_n)}$. There is a GKM function $G \in R$ such that h coincides with $f^{(0)}{}^{I_{n-1}}_{i'_n} \prod_{0 \leq l < k'} (\tau_{n+l} + t_{i'_n}) G$ on $W^{(I_{n-1}, n+k')}_{(I'_{n-1}, -i'_n)}$.*

We show this claim by induction on k'. For $w \in W^{(I_{n-1}, n+k')}_{(I'_{n-1}, -i'_n)}$, by the GKM condition, $h(w)$ belongs to the ideal generated by the following elements of $H^*(\mathcal{F}_4)$.

$$\begin{aligned} w(t_{n+l} - t_{n+k'}) &= w(t_{n+l}) + t_{i'_n} && \text{for } 0 \leq l < k' \\ w(t_{n+l} + t_{n+k'}) &= w(t_{n+l}) - t_{i'_n} && \text{for } 0 \leq l \leq 4-n,\ l \neq k' \\ w(t_{n+k'}) &= -t_{i'_n} \end{aligned}$$

For $w \in W^{(I_{n-1},n+k')}_{(I'_{n-1},-i'_n)}$, by the equation (5.2), there is an element $g_w \in H^*(BT)$ such that

$$h(w) = f^{(0)}{}^{I_{n-1}}_{i'_n}(w) \prod_{0\leq l<k'} (\tau_{n+l} + t_{i'_n})(w) g_w.$$

One can verify that a function G' given by $G'(w) = g_w$ satisfies the GKM condition on $W^{(I_{n-1},n+k')}_{(I'_{n-1},-i'_n)}$ similarly as above. By (decreasing) induction on n there is a GKM function $G \in R$ such that G and G' coincide on $W^{(I_{n-1},n+k')}_{(I'_{n-1},-i'_n)}$. Then

$$h - f^{(0)}{}^{I_{n-1}}_{i'_n} \prod_{0\leq l<k'} (\tau_{n+l} + t_{i'_n}) G = 0$$

$$\text{on } \coprod_{0\leq l\leq 4-n} W^{(I_{n-1},n+l)}_{(I'_{n-1},i'_n)} \sqcup \coprod_{0\leq l\leq k'} W^{(I_{n-1},n+l)}_{(I'_{n-1},-i'_n)}.$$

Therefore the induction on k' proceeds. By Proposition 5.1 the function

$$f^{(0)}{}^{I_{n-1}}_{i'_n} = \frac{1}{2} \prod_{0\leq l\leq 4-n} (\tau_{n+l} - t_{i'_n})$$

can be replaced by a polynomial in γ_j's ($1 \leq j \leq 4$) over $H^*(BT)$. Therefore the (decreasing) induction on n proceeds, and we may assume h vanishes on $W(Spin(9)) = W \sqcup \kappa W$.

Next we show that for a GKM function h which vanishes on $W(Spin(9))$, there is a GKM function $G \in R$ such that $h - \omega G = 0$ on $W(Spin(9)) \sqcup \rho W(Spin(9))$, where $\omega = e_4(\tau) - e_4(t) - 2\gamma_4$ vanishes on $W(Spin(9))$. Then the GKM condition says that for $w \in \rho W(Spin(9))$, $h(w)$ belongs to the ideal $(\rho^2 t_i \mid i = 1, 2, 3, 4)$. For $w \in \rho W(Spin(9))$, there is an element $g_w \in H^*(BT)$ such that

$$h(w) = -e_4(\rho^2 t) g_w = \omega(w) g_w.$$

It is obvious that a function G' given by $G'(w) = g_w$ satisfies the GKM condition on $\rho W(Spin(9))$, since the edges in the GKM subgraph induced by $\rho W(Spin(9))$ have the long roots or ρt_i as their labels and the all roots of F_4 are prime in $H^*(BT)$. Then there is a GKM function G such that $G = G'$ on $\rho W(Spin(9))$. It is proved similarly as above, changing $W^{I_n}_{I'_n}$ to $\rho W^{I_n}_{I'_n}$, $\tau_{k_s} - t_{i'_n}$ to $\tau_{k_s} - \rho t_{i'_n}$, and $f^{(0)}{}^{I_{n-1}}_{i'_n}$ to $f^{(1)}{}^{I_{n-1}}_{i'_n}$.

Finally we show that for a GKM function h which vanishes on $W(Spin(9)) \sqcup \rho W(Spin(9))$, there is a GKM function $G \in R$ such that $h - \omega(\omega + e_4(\rho^2 t))G = 0$ as a GKM function, where $\omega + e_4(\rho^2 t)$ vanishes on $\rho W(Spin(9))$. It is proved similarly as above that for $w \in \rho^2 W(Spin(9))$, $h(w)$ belongs to the ideal $(t_i, \rho t_i \mid i = 1, 2, 3, 4)$. For $w \in \rho^2 W(Spin(9))$, there is an element $g_w \in H^*(BT)$ such that

$$h(w) = -e_4(\rho t) e_4(t) g_w = \omega(w)(\omega(w) + e_4(\rho^2 t)) g_w,$$

where the latter equality is due to the relation (4.4). Then a function G' given by $G'(w) = g_w$ satisfies the GKM condition on $\rho^2 W(Spin(9))$, and there is a GKM function G such that $G = G'$ on $\rho^2 W(Spin(9))$. It is proved similarly as above, changing $W^{I_n}_{I'_n}$ to $\rho^2 W^{I_n}_{I'_n}$, $\tau_{k_s} - t_{i'_n}$ to $\tau_{k_s} - \rho^2 t_{i'_n}$, and $f^{(0)}{}^{I_{n-1}}_{i'_n}$ to $f^{(2)}{}^{I_{n-1}}_{i'_n}$. The proof is completed. $\square$

6 Proof of Proposition 4.2

Let us prove Proposition 4.2 and that the relations (4.13), (4.14), (4.15), and (4.16) hold. For convenience we call the relations (4.5), (4.6), (4.7), (4.8), (4.9), (4.10), (4.11), (4.12), (4.13), (4.14), (4.15), and (4.16) by q_1', q_1, q_2, q_3, q_4, q_6, q_8, q_{12}, p_2', p_2, p_4, and p_{12}, respectively.

Proof of Proposition 4.2. The relations q_1', q_1, q_3, and p_2' hold obviously by definition, and the relation p_{12} and q_{12} hold by the equation (4.2) and (4.3). To show p_2, p_4, q_6, and q_8, we claim that the following relations hold in $H_T^*(\mathcal{F}_4)$.

$$e_1(\tau^2) - e_1(t^2) = 0 \tag{6.1}$$

$$e_2(\tau^2) - e_2(t^2) - 6\omega = 0 \tag{6.2}$$

$$e_3(\tau^2) - e_3(t^2) - e_1(t^2)u = 0 \tag{6.3}$$

$$e_4(\tau^2) - e_4(t^2) + 3\omega^2 - 2(e_4(\rho t) - e_4(\rho^2 t))\omega = 0 \tag{6.4}$$

The left-hand side functions of these equations are constant on each $\rho^\varepsilon W(Spin(9))$ for $\varepsilon = 0, 1, 2$. Calculations of each values on $\rho^\varepsilon W(Spin(9))$ with the equation (4.4) show (6.1), (6.2), (6.3), and (6.4) hold.

We show that (6.1), (6.2), (6.3), and (6.4) are divisible by 4 to deduce p_2, p_4, q_6, and q_8. Let $X = -6ux^4 + e_1(t^2)ux^6 - (3u^2 + 2(e_4(\rho t) - e_4(\rho^2 t))u)x^8$. It follows from (6.1), (6.2), (6.3), and (6.4) that

$$\begin{aligned}
0 &= \prod_{i=1}^{4}(1 - \tau_i^2 x^2) - \prod_{i=1}^{4}(1 - t_i^2 x^2) + X \\
&= \sum_{k=0}^{4}(1 + (-1)^k e_k(\tau)x^k)\sum_{k=0}^{4}(1 + e_k(\tau)x^k) - \sum_{k=0}^{4}(1 + (-1)^k e_k(t)x^k)\sum_{k=0}^{4}(1 + e_k(t)x^k) + X.
\end{aligned}$$

We modify the right-hand side by the relation q_1, p_2', and q_3 to obtain

$$\begin{aligned}
&4\sum_{k=1}^{3}(-1)^k\gamma_k^2 x^{2k} - 8\gamma_1\gamma_3 x^4 + 4\sum_{k=1}^{3}\sum_{i=n_k}^{m_k}(-1)^i\gamma_i e_{2k-i}(t)x^{2k} \\
&\quad + 2(e_4(\tau) - e_4(t))x^4 + 4\gamma_2 e_4(\tau)x^6 + 2e_2(t)(e_4(\tau) - e_4(t))x^6 + (e_4(\tau)^2 - e_4(t)^2)x^8 + X,
\end{aligned}$$

where $n_k = \max\{1, 2k-3\}$ and $m_k = \min\{3, 2k\}$. This calculation is similar to the calculation in [FIM, Proof of Lemma 5.5]. However remark that $\gamma_4 \neq \frac{1}{2}(e_4(\tau) - e_4(t))$. Then comparing the coefficients we obtain

$$\begin{aligned}
0 &= -4\gamma_1^2 + 4(-\gamma_1 e_1(t) + \gamma_2), \\
0 &= 4\gamma_2^2 - 8\gamma_1\gamma_3 + 4(-\gamma_1 e_3(t) + \gamma_2 e_2(t) - \gamma_3 e_1(t)) + 2(e_4(\tau) - e_4(t)) - 6\omega \\
&= 4\gamma_2^2 - 4(e_1(\tau) - e_1(t))\gamma_3 + 4(-\gamma_1 e_3(t) + \gamma_2 e_2(t) - \gamma_3 e_1(t)) - 4(e_4(\tau) - e_4(t)) + 12\gamma_4, \\
0 &= -4\gamma_3^2 - 4\gamma_3 e_3(t) + 4\gamma_2 e_4(\tau) + 2e_2(t)(e_4(\tau) - e_4(t)) + e_1(t^2)\omega \\
&= -4\gamma_3^2 - 4\gamma_3 e_3(t) + 4\gamma_2 e_4(\tau) + 2e_2(t)(\omega + 2\gamma_4) + (e_1(t)^2 - 2e_2(t))\omega \\
&= -4\gamma_3^2 - 4\gamma_3 e_3(t) + 4\gamma_2 e_4(\tau) + 4e_2(t)\gamma_4 + 4\gamma^2\omega.
\end{aligned}$$

Regarding GKM functions as elements of $\mathrm{Map}(W(G), H^*(BT)\otimes\mathbb{Q})$, we can divide them by 4 to obtain

$$\begin{aligned}
\gamma_1^2 &= -\gamma_1 e_1(t) + \gamma_2,\\
\gamma_2^2 &= e_1(\tau)\gamma_3 + \gamma_1 e_3(t) - \gamma_2 e_2(t) + e_4(\tau) - e_4(t) - 3\gamma_4,\\
\gamma_3^2 &= -\gamma_3 e_3(t) + \gamma_2 e_4(\tau) + \gamma_4 e_2(t) + \gamma^2\omega.
\end{aligned}$$

Since each sides of these equations remain polynomials in $H^*(BT)$-valued GKM functions over $\mathbb{Z}$, these equations hold in $H^*(\mathcal{F}_4) \subset \mathrm{Map}(W(G), H^*(BT))$. The first and the second equations show the relations p_2 and p_4 respectively. The third equation with p_2 shows the relation q_6. Obviously the relations p_2 and q_1 show the relation q_2. We modify the left-hand side of the equation (6.4) by $\omega = e_4(\tau) - e_4(t) - 2\gamma_4$ to obtain

$$\begin{aligned}
&4e_4(\tau)^2 + (-12\gamma_4 - 6e_4(t) - 2e_4(\rho t) + 2e_4(\rho^2 t))e_4(\tau) + 12\gamma_4^2\\
&+ (12e_4(t) + 4e_4(\rho t) - 4e_4(\rho^2 t))\gamma_4 + 2e_4(t)^2 + (2e_4(\rho t) - 2e_4(\rho^2 t))e_4(t)
\end{aligned}$$

Since $e_4(t^2) = e_4(t)^2$, by the equation (4.4) it is equal to

$$4e_4(\tau)^2+(-12\gamma_4-4e_4(t)+4e_4(\rho^2t))e_4(\tau)+12\gamma_4^2+(12e_4(t)+4e_4(\rho t)-4e_4(\rho^2t))\gamma_4-4e_4(\rho^2t)e_4(t).$$

This is divided by 4 to yield

$$e_4(\tau)^2 + 3\gamma_4(\gamma_4 - e_4(\tau) + e_4(t)) - \gamma_4 e_4(t) - e_4(\tau)e_4(t) + e_4(\rho^2 t)(e_4(\tau) - e_4(t) - 2\gamma_4).$$

Erasing γ_4 by the relations p_4 and (4.17) we obtain the relation q_8 as

$$\begin{aligned}
&e_4(\tau)^2 - A^2 + (e_4(\tau) + 2\omega)A + 2\omega(e_4(t) - e_4(\tau)) - \omega e_4(t) - e_4(\tau)e_4(t) + e_4(\rho^2 t)\omega\\
= &e_4(\tau)^2 - A^2 + (e_4(\tau) + 2\omega)A - e_4(\tau)(2\omega + e_4(t)) - \omega e_4(\rho t).
\end{aligned}$$

Finally the relation p_4 with p_2 and (4.17) shows the relation q_4. $\square$

7 Proof of Lemma 4.2

We will prove Lemma 4.2 by the argument of regular sequences. The following theorems and propositions are useful. Proposition 7.1 and 7.2 are obvious by definition.

Theorem 7.1 ([M, Theorem 16.1]). *If $a_1, \dots, a_n$ is a regular sequence, then so is $a_1^{v_1}, \dots, a_n^{v_n}$ for any positive integers $v_1, \dots, v_n$.*

Theorem 7.2 ([M, Corollary of Theorem 16.3]). *Let A be a Noetherian ring and $\mathbb{N}$-graded. If $a_1, \dots, a_n$ is a regular sequence in A and each a_i is homogeneous of positive degree, then any permutation of $a_1, \dots, a_n$ is again a regular sequence.*

Proposition 7.1. *If $a_1, \dots, a_n$ is a regular sequence, then so is $a_1, \dots, a_{i-1}, a_i + b, a_{i+1}, \dots, a_n$ for $1 \le i \le n$ and any $b \in (a_1, \dots, a_{i-1})$.*

Proposition 7.2. *If $a_1, \dots, a_n$ is a regular sequence, then so is $a_1, \dots, a_{i-1}, a_{i+1}, \dots, a_n$ for $1 \leq i \leq n$.*

Proof of Lemma 4.2. Let R denote the polynomial ring $\mathbb{Z}[t_i, \gamma, \tau_i, \gamma_1, \gamma_3, \omega \mid 1 \leq i \leq 4]$ and I denote the ideal generated by r_1', r_1, r_2, r_3, r_4, r_6, r_8, and r_{12} in R. We will show the sequence

$$r_1', r_{12}, r_4, r_3, r_2, r_1, r_6, r_8$$

is a regular sequence in $R \otimes \mathbb{Z}/2\mathbb{Z}$ and the sequence

$$r_1', r_{12}, r_8, r_6, r_4, r_2, r_3, r_1$$

is a regular sequence in $R \otimes \mathbb{Z}/p\mathbb{Z}$ for any prime $p \geq 3$. If these sequences are regular, since $\mathbb{Z}/p\mathbb{Z}$ is a field, by [NS, Theorem 5.5.1] the Poincaré series of $(R/I) \otimes \mathbb{Z}/p\mathbb{Z}$ is calculated from the degrees of the generators and the relations as

$$\begin{aligned}
&\frac{(1-x^2)(\prod_{j=1}^{3}(1-x^{2j}))(1-x^{24})\prod_{i=1}^{4}(1-x^{4i})}{(1-x^2)^4(1-x^2)(1-x^2)^4\prod_{i=1}^{4}(1-x^{2i})} \\
&= \frac{1}{(1-x^2)^4}(1+x^8+x^{16})\prod_{i=1}^{4}\frac{1-x^{4i}}{1-x^2}
\end{aligned}$$

for any prime p. Then R/I is free as a $\mathbb{Z}$-module and its Poincaré series coincides with that of $H_T^*(F_4/T)$ by Proposition 4.1, completing the proof.

When $p = 2$, we have

$$\begin{aligned}
r_1' &= e_1(t) \\
r_1 &= -(e_1(\tau) - e_1(t)) \\
r_2 &= -(e_2(\tau) - e_2(t)) + (e_1(\tau) - e_1(t))e_1(t) \\
r_3 &= -(e_3(\tau) - e_3(t)) \\
r_4 &= e_4(\tau) - e_4(t) - 3\omega
\end{aligned}$$

in $(R/I) \otimes \mathbb{Z}/2\mathbb{Z}$. Especially we have

$$A \equiv \gamma_1^4 - \gamma_3 e_1(\tau) \qquad \text{mod } (e_i(t) \mid 1 \leq i \leq 4) \tag{7.1}$$

$$r_6 \equiv \gamma_3^2 \qquad \text{mod } (\gamma, e_i(t), e_4(\tau) \mid 1 \leq i \leq 4), \tag{7.2}$$

and by the equation (7.1)

$$r_8 \equiv -\gamma_1^8 \quad \text{mod } (e_i(t), e_4(\tau), \omega \mid 1 \leq i \leq 4).$$

It is well known that the sequence of the elementary symmetric polynomials

$$e_1(x), e_2(x), \dots, e_n(x),$$

that is, the sequence of the Chern classes is regular in $\mathbb{Z}/p\mathbb{Z}[x_i \mid 1 \leq i \leq n]$ for any prime p. Since the polynomial ring R is Noetherian, by Theorem 7.2 the sequence

$$\gamma, e_1(t), e_2(t), e_3(t), e_4(t), \omega, e_4(\tau), e_3(\tau), e_2(\tau), e_1(\tau), \gamma_3^2, -\gamma_1^8$$

is regular in R. We modify this sequence by Theorem 7.1 and Proposition 7.1 to obtain the following regular sequence.

$$\gamma, r'_1, e_2(t), e_3(t), e_4(t), \omega^3, r_4, r_3, r_2, r_1, r_6, r_8$$

Since $\rho^2 t_4 = -\gamma$ and $e_4(\rho t) = -e_4(t) - e_4(\rho^2 t) \equiv 0 \bmod (\gamma, e_4(t))$, by Proposition 7.1

$$\gamma, r'_1, e_2(t), e_3(t), e_4(t), r_{12}, r_4, r_3, r_2, r_1, r_6, r_8$$

is a regular sequence. Hence

$$r'_1, r_{12}, r_4, r_3, r_2, r_1, r_6, r_8$$

is a regular sequence by Proposition 7.2.

When $p \geq 3$, we will show the regularity of the sequence $r'_1, r_{12}, r_8, r_6, r_4, r_2, r_3, r_1$ by the modification as follows. Put $r'_2 = \gamma_2 - \gamma_1^4 - \gamma_1 e_1(t)$ and $r'_4 = \gamma_4 - \omega + A$. They correspond the relations (4.14) and (4.17). Then the following ring isomorphism holds.

$$\begin{aligned} &(\mathbb{Z}[t_i, \gamma, \tau_i, \gamma_1, \gamma_3, \omega \mid 1 \leq i \leq 4]/(r'_1, r_1, r_2, r_3, r_4, r_6, r_8, r_{12}) \otimes \mathbb{Z}/p\mathbb{Z} \\ \cong\ &(\mathbb{Z}[t_i, \gamma, \tau_i, \gamma_i, \omega \mid 1 \leq i \leq 4]/(r'_1, r_1, r_2, r_3, r_4, r_6, r_8, r_{12}, r'_2, r'_4) \otimes \mathbb{Z}/p\mathbb{Z} \end{aligned} \tag{7.3}$$

Let e_1, e_2, e_3, and e_4 be the left-hand sides of (6.1), (6.2), (6.3), and (6.4) respectively, namely

$$\begin{aligned} e_1 &= e_1(\tau^2) - e_2(t^2), & e_2 &= e_2(\tau^2) - e_2(t^2), \\ e_3 &= e_3(\tau^2) - e_2(t^2) - e_1(t^2)\omega, & e_4 &= e_4(\tau^2) - e_4(t^2) + (e_4(\rho t) - e_4(\rho^2 t))\omega. \end{aligned}$$

Recall that e_1, e_2, e_3, and e_4 are divided by 4 to yield r_2, r_4, r_6, and r_8 respectively. Since $p \geq 3$, 2 is invertible in $\mathbb{Z}/p\mathbb{Z}$. Then (7.3) is isomorphic to

$$\begin{aligned} &(\mathbb{Z}[t_i, \gamma, \tau_i, \gamma_i, \omega \mid 1 \leq i \leq 4]/(r'_1, r_1, e_2, r_3, e_4, e_6, e_8, r_{12}, r'_2, r'_4) \otimes \mathbb{Z}/p\mathbb{Z} \\ \cong\ &(\mathbb{Z}[t_i, \gamma, \tau_i, \gamma_1, \gamma_3, \omega \mid 1 \leq i \leq 4]/(r'_1, r_1, e_2, r_3, e_4, e_6, e_8, r_{12}) \otimes \mathbb{Z}/p\mathbb{Z}. \end{aligned}$$

Since γ_1 and γ_3 can be canceled by the relations $r_1 = 0$ and $r_3 = 0$, it is isomorphic to

$$(\mathbb{Z}[t_i, \gamma, \tau_i, \omega \mid 1 \leq i \leq 4]/(r'_1, e_2, e_4, e_6, e_8, r_{12}) \otimes \mathbb{Z}/p\mathbb{Z}.$$

Hence it is sufficient to show that the sequence $r'_1, r_{12}, e_8, e_6, e_4, e_2$ is regular in $R \otimes \mathbb{Z}/p\mathbb{Z}$. It is well known that the sequence of elementary symmetric polynomial in $\{x_i^2\}_{i=1}^n$

$$e_1(x^2), e_2(x^2), \ldots, e_n(x^2)$$

that is, the sequence of the Pontryagin classes is regular in $\mathbb{Z}/p\mathbb{Z}[x_i \mid 1 \leq i \leq n]$ for any prime p. Since the polynomial ring R is Noetherian, by Theorem 7.2 the sequence

$$\gamma, e_1(t), e_2(t), e_3(t), e_4(t), \omega, e_4(\tau^2), e_3(\tau^2), e_2(\tau^2), e_1(\tau^2)$$

is regular in R. We modify this sequence by Theorem 7.1 and Proposition 7.1 to obtain the following regular sequence.

$$\gamma, r'_1, e_2(t), e_3(t), e_4(t), r_{12}, e_8, e_6, e_4, e_2$$

Hence

$$r'_1, r_{12}, e_8, e_6, e_4, e_2$$

is a regular sequence by Proposition 7.2. Therefore R/I is free as a $\mathbb{Z}$-module and its Poincaré series coincides with that of $H_T^*(F_4/T)$. $\square$

8 Proof of Corollary 1.1

Proof. Let R be the polynomial ring $\mathbb{Z}[\tau_i, \gamma_1, \gamma_3, \omega \mid 1 \le i \le 4]$. By the argument in Section 1 we have the isomorphisms

$$\begin{aligned} H^*(F_4/T) &\cong H^*_T(F_4/T)/(t_1, t_2, t_3, t_4, \gamma) \\ &\cong R/(\overline{r}_1, \overline{r}_2, \overline{r}_3, \overline{r}'_4, \overline{r}_6, \overline{r}'_8, \overline{r}_{12}), \end{aligned}$$

where

$$\begin{aligned} \overline{r}'_4 &= e_4(\tau) - 2\gamma_3 e_1(\tau) + 2\gamma_1^4 - 3\omega, \\ \overline{r}'_8 &= e_4(\tau)^2 - (\gamma_1^4 - 2\gamma_3 e_1(\tau))^2 + (e_4(\tau) + 2\omega)(\gamma_1^4 - 2\gamma_3 e_1(\tau)) - 2\omega e_4(\tau). \end{aligned}$$

We show

$$\overline{r}_4 \equiv \overline{r}'_4 \mod (\overline{r}_1, \overline{r}_3), \tag{8.1}$$
$$\overline{r}_8 \equiv \overline{r}'_8 \mod (\overline{r}_1, \overline{r}_3, \overline{r}_4, \overline{r}_6) \tag{8.2}$$

to prove the equality of the two ideals $(\overline{r}_1, \overline{r}_2, \overline{r}_3, \overline{r}'_4, \overline{r}_6, \overline{r}'_8, \overline{r}_{12})$ and $(\overline{r}_1, \overline{r}_2, \overline{r}_3, \overline{r}_4, \overline{r}_6, \overline{r}_8, \overline{r}_{12})$. Since the relation $\overline{r}_1 = \overline{r}_3 = 0$ means $e_i(\tau) = 2\gamma_i$ for $i = 1, 3$, the equation (8.1) holds. By the relations $\overline{r}_4 = 0$ we can deform $\overline{r}_8$ as

$$3e_4(\tau)\gamma_1^4 - \gamma_1^8 + (e_4(\tau) - 4\gamma_1\gamma_3 + 2\gamma_1^4)(\omega + 2\gamma_1\gamma_3) \tag{8.3}$$

in $R/(\overline{r}_1, \overline{r}_3, \overline{r}_4)$. On the other hand $\overline{r}'_8$ yields

$$e_4(\tau)^2 - (\gamma_1^4 - 2\gamma_1\gamma_3)^2 + (e_4(\tau) + 2\omega)(\gamma_1^4 - 2\gamma_1\gamma_3) - 2\omega e_4(\tau) \tag{8.4}$$

in $R/(\overline{r}_1, \overline{r}_3, \overline{r}_4)$. Then the difference of (8.4) and (8.3) is

$$\begin{aligned} & e_4(\tau)^2 + 4\gamma_1^2\gamma_3^2 - 2\gamma_1^4 e_4(\tau) - 4\gamma_1\gamma_3 e_4(\tau) - 3\omega e_4(\tau) \\ &= (e_4(\tau) + 2\gamma_1^4 - 4\gamma_1\gamma_3 - 3\omega)e_4(\tau) + 4\gamma_1^2(-\gamma_1^2 e_4(\tau) + \gamma_3^2) \end{aligned} \tag{8.5}$$

By the relations $\overline{r}_4 = 0$ and $\overline{r}_6 = 0$, (8.5) equals to zero in $R/(\overline{r}_1, \overline{r}_3, \overline{r}_4, \overline{r}_6)$, and then (8.2) holds. $\square$

References

[A] J. F. Adams, *Lectures on exceptional Lie groups* University of Chicago Press, Chicago, 1996.

[F] Y. Fukukawa, *The cohomology ring of the GKM graph of a flag manifold of type G_2*, arXiv:1207.5229v1.

[FIM] Y. Fukukawa, H. Ishida, and M. Masuda, *The cohomology ring of the GKM graph of a flag manifold of classical type*, arXiv:1104.1832v1.

[GKM] M. Goresky, R. Kottwitz, and R. MacPherson, *Equivariant cohomology, Koszul duality and the localization theorem*, Invent. Math. **131** (1998), no. 1, 25-83.

[GZ] V. Guillemin and C. Zara, *1-skeleta, Betti numbers, and equivariant cohomology*, Duke Math. J. **107** (2001), no. 2, 283-349.

[HHH] M. Harada, A. Henriques, and T. Holm, *Computation of generalized equivariant cohomologies of Kac-Moody flag varieties*, Adv. Math. **197** (2005), no 1, 198-221.

[M] H. Matsumura, *Commutative ring theory*, Cambridge studies in advanced mathematics, **8**, Cambridge University Press, Cambridge, 1986.

[NS] M. D. Neusel, L. Smith, *Invariant theory of finite groups*, Mathematical Surveys and Monographs Volume **94**, American Mathematical Society, Providence, 2002.

[TW] H. Toda and T. Watanabe, *The integral cohomology ring of F_4/T and E_6/T*, J. Math. Kyoto Univ. **14-2** (1974), 257-286.